

AD-753 964

EQUATION OF THE STATE OF A HELIUM AND
NITROGEN MIXTURE

V. N. Popov, et al

Foreign Technology Division
Wright-Patterson Air Force Base, Ohio

27 October 1972

DISTRIBUTED BY:

NTIS

National Technical Information Service
U. S. DEPARTMENT OF COMMERCE
5285 Port Royal Road, Springfield Va. 22151

AD753964

FTD-HT-23-956-72



FOREIGN TECHNOLOGY DIVISION



EQUATION OF THE STATE OF A HELIUM AND
NITROGEN MIXTURE

by

V. N. Popov, V. I. Chernyshov



Approved for public release;
distribution unlimited.

Reproduced by
**NATIONAL TECHNICAL
INFORMATION SERVICE**
U S Department of Commerce
Springfield VA 22151

UNCLASSIFIED

Security Classification

DOCUMENT CONTROL DATA - R & D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author) Foreign Technology Division Air Force Systems Command U. S. Air Force		2a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED	
		2b. GROUP	
3. REPORT TITLE EQUATION OF THE STATE OF A HELIUM AND NITROGEN MIXTURE			
4. DESCRIPTIVE NOTES (Type of report and inclusive dates) Translation			
5. AUTHOR(S) (First name, middle initial, last name) Popov, V.N.; Chernvshov, V.I.			
6. REPORT DATE 1971	7a. TOTAL NO. OF PAGES 13	7b. NO. OF REFS 18	
8a. CONTRACT OR GRANT NO. a. PROJECT NO. 7343 b. c.		8b. ORIGINATOR'S REPORT NUMBER(S) FTD-HT-23-956-72	
		8c. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
9. DISTRIBUTION STATEMENT Approved for public release; distribution unlimited.			
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY Foreign Technology Division Wright-Patterson AFB, Ohio	
13. ABSTRACT The equation of state describing the experimental data within the temperature range of 273.15 to 773.15°K and a pressure up to 400 bars was obtained as a result of processing experimental data with respect to the compressibility of helium, nitrogen, and mixtures of them on a digital computer. Comparison of the calculated values of the coefficients of compressibility and density of the mixture of helium with nitrogen calculated by the proposed equation of state with the experimental values demonstrates satisfactory comparison in the entire temperature and pressure range encompassed by the experiment. AP1207329 1A			

DD FORM 1 NOV 61 1473

UNCLASSIFIED

Security Classification

Inventory Classification

14

KEY WORDS

Equation of State
Helium
Nitrogen
Digital Computer
Gas Density
Compressible Gas

ib

UNCLASSIFIED

Security Classification

EDITED TRANSLATION

FTD-HT-23-956-72

EQUATION OF THE STATE OF A HELIUM AND NITROGEN
MIXTURE

By: V. N. Popov, V. I. Chernyshov

English pages: 8

Source: AN SSSR. Izvestiya. Energetika i
Transport, No. 3, 1971, pp. 162-165.

Requester: ASD

Translated by: Joseph E. Pearson

Approved for public release;
distribution unlimited.

THIS TRANSLATION IS A RENDITION OF THE ORIGINAL FOREIGN TEXT WITHOUT ANY ANALYTICAL OR EDITORIAL COMMENT. STATEMENTS OR THEORIES ADVOCATED OR IMPLIED ARE THOSE OF THE SOURCE AND DO NOT NECESSARILY REFLECT THE POSITION OR OPINION OF THE FOREIGN TECHNOLOGY DIVISION.

PREPARED BY:

TRANSLATION DIVISION
FOREIGN TECHNOLOGY DIVISION
WP-afb, OHIO.

U. S. BOARD ON GEOGRAPHIC NAMES TRANSLITERATION SYSTEM

Block	Italic	Transliteration	Block	Italic	Transliteration
А а	<i>А а</i>	A, a	Р р	<i>Р р</i>	R, r
Б б	<i>Б б</i>	B, b	С с	<i>С с</i>	S, s
В в	<i>В в</i>	V, v	Т т	<i>Т т</i>	T, t
Г г	<i>Г г</i>	G, g	У у	<i>У у</i>	U, u
Д д	<i>Д д</i>	D, d	Ф ф	<i>Ф ф</i>	F, f
Е е	<i>Е е</i>	Ye, ye; E, e*	Х х	<i>Х х</i>	Kh, kh
Ж ж	<i>Ж ж</i>	Zh, zh	Ц ц	<i>Ц ц</i>	Ts, ts
З з	<i>З з</i>	Z, z	Ч ч	<i>Ч ч</i>	Ch, ch
И и	<i>И и</i>	I, i	Ш ш	<i>Ш ш</i>	Sh, sh
Й й	<i>Й й</i>	Y, y	Щ щ	<i>Щ щ</i>	Shch, shch
К к	<i>К к</i>	K, k	Ъ ъ	<i>Ъ ъ</i>	"
Л л	<i>Л л</i>	L, l	Ы ы	<i>Ы ы</i>	Y, y
М м	<i>М м</i>	M, m	Ь ь	<i>Ь ь</i>	'
Н н	<i>Н н</i>	N, n	Э э	<i>Э э</i>	E, e
О о	<i>О о</i>	O, o	Ю ю	<i>Ю ю</i>	Yu, yu
П п	<i>П п</i>	P, p	Я я	<i>Я я</i>	Ya, ya

* ye initially, after vowels, and after ъ, ы; e elsewhere.
 When written as ѣ in Russian, transliterate as yě or ѣ.
 The use of diacritical marks is preferred, but such marks
 may be omitted when expediency dictates.

EQUATION OF THE STATE OF A HELIUM AND NITROGEN MIXTURE

V. N. Popov, V. I. Chernyshov
(Moscow)

As a result of the digital computer processing of the experimental data on the compressibility of helium, nitrogen and their mixtures, an equation of state which describes the experimental data in the 273.15-773.15°K temperature interval and under a pressure of 400 bar, was obtained.

A comparison of the calculated values of the coefficients of compressibility and density of a mixture of helium and nitrogen, calculated by the proposed equation of state, with the experimental data shows a satisfactory coincidence in the entire temperature and pressure region, encompassed by the experiment. Tables 5, Bibliographical entries 18, Pages 162-165.

A mixture of helium and nitrogen is employed in technology as a heat-transfer agent, thus for carrying out hydrodynamic calculations and solving many heat-exchange problems it is necessary to have reliable data available on the thermal, calorific properties, and also on the transfer properties of this mixture in a broad range of variation of the parameters of state.

In the present article by the digital computer processing of the experimental data [1-4] for helium, nitrogen, their mixtures, equations of state were obtained, by which detailed tables of the values of the coefficients of compressibility and density were computed in the temperature range from 273.15 up to 773.15°K and in the pressure range from 50 to 400 bar.

Equation of the State of Helium. For the processing of all the experimental data on helium in the temperature and pressure intervals examined by us, an equation of state in the form of a polynomial for degrees of density was employed

$$z = 1 + B_{11}\rho + C_{111}\rho^2. \quad (1)$$

The dependences $B_{11} = f(T)$ and $C_{111} = f(T)$ were approximated by a polynomial for the degrees $\tau = 1000/T$

$$B_{11}(T) = B_{11}^0(T) + \sum_{i=1}^2 b_i \tau^i, \quad (2)$$

$$C_{111}(T) = \sum_{i=1}^2 c_i \tau^i. \quad (3)$$

The coefficients of these dependences were determined on a digital computer by the method of least squares in two variables for the whole aggregate of available experimental data for helium [5-11], also including our experimental data [3]. The values of coefficients b_i and c_i are given in Table 1. Dependence $B_{11}^0(T)$ is taken in accordance with [12] and has the form:

$$B_{11}^0(T) = \left(0.001801 + \frac{0.02169\tau}{1 + 1.89\tau} \right). \quad (4)$$

A total of 254 experimental points were taken for processing. The deviations of the calculated values of the coefficient of compressibility from the experimental values for 188 points were less than $\pm 0.05\%$, for 47 points less than $\pm 0.10\%$ and for 19 points less than $\pm 0.25\%$.

Table 1. The values of the coefficients of polynomials (2) and (3) for helium.

$b_i \cdot 10^4,$ $m^3/kmole$	b_0	+0.45730
	b_1	+1.41850
	b_2	-0.36636
$c_i \cdot 10^6,$ $m^6/(kmole)^2$	c_0	+0.34964
	c_1	+6.58490
	c_2	-0.90130

The equation of state of nitrogen. In composing the equation of state for nitrogen we used experimental data [13-15], which encompasses the temperature and pressure intervals examined by us and also our experimental data [2, 4]. The equation of state was composed in two stages.

In the first stage, the graphico-analytical method was used, which was examined as a first approximation. For approximating the dependence of the second virial coefficient $B_{22}^0(T)$ on temperature for the whole aggregate of experimental data [7, 13, 15, 16] with respect to $B_{22}^0(T)$ a polynomial for degrees of τ was selected

$$B_{22}^0(T) = \sum_{i=0}^4 b_i / \tau^i. \quad (5)$$

The values of the coefficients of b_i are given in Table 2. In the second stage the final equation of state is used in the form

$$z = 1 + B_{22}\rho + C_{222}\rho^2 + D_2\rho^3 + E_2\rho^4. \quad (6)$$

Dependences $B_{22}(T)$, $C_{222}(T)$, $D_2(T)$ and $E_2(T)$ were approximated by a polynomial for degrees $\tau = 1000/T$

$$B_{22}(T) = B_{22}^0(T),$$

$$C_{222}(T) = \sum_{i=0}^4 c_i \tau^i, \quad D_2(T) = \sum_{i=0}^4 d_i \tau^i, \quad E_2(T) = \sum_{i=0}^4 e_i \tau^i. \quad (7)$$

The coefficients of these dependences (7) were determined on a digital computer by the method of least squares for the whole aggregate of available experimental points for nitrogen. The values of the coefficients of c_i , d_i , e_i are given in Table 3.

Table 2. The values of the coefficients b_i of polynomial (5) for nitrogen.

$b_i \cdot 10^5$	b_0	-0.14873
m^3/kmole	b_1	+0.91396
	b_2	-2.23460
	b_3	+2.05920
	b_4	-0.82300

A total of 140 experimental points were taken for processing. The deviations of the calculated values of the coefficient of compressibility from the experimental values for 114 points was less than $\pm 0.05\%$, for 17 points less than $\pm 0.10\%$ and for 9 points less than $\pm 0.25\%$.

The equation of state of a mixture of helium and nitrogen. For composing the equation of state of a mixture in virial form the processing of the experimental data was carried out in two stages. In the first stage the graphico-analytical method of processing was used, which was examined as a first approximation. This method is described in detail in work [2].

For the final processing of the experimental data [2, 4, 15] in the temperature range 273.15-773.15°K and pressure range 50-400 bar an equation in the form of a polynomial for degrees of density was selected

$$z_{CM} = 1 + B_{CM}\rho + C_{CM}\rho^2 + D_{CM}\rho^3 + E_{CM}\rho^4. \quad (8)$$

Functions B_{CM} , C_{CM} , D_{CM} , E_{CM} , have the following form:

$$\begin{aligned} B_{CM}(T, x) &= B_{11}x_1^2 + 2B_{12}x_1x_2 + B_{22}x_2^2, \\ C_{CM}(T, x) &= C_{111}x_1^3 + 3C_{112}x_1^2x_2 + 3C_{122}x_1x_2^2 + C_{222}x_2^3, \\ D_{CM}(T, x) &= \gamma_1(1-x_1) + (D_2 - \gamma_1)(1-x_1)^2 + \gamma_2(1-x_1)^2x_1, \\ E_{CM}(T, x) &= a_1(1-x_1) + (E_2 - a_1)(1-x_1)^2. \end{aligned} \quad (9)$$

Table 3. The values of the coefficient of the polynomials (7) in the equation of state for nitrogen.

$c_i \cdot 10^2$	c_0	-0.62141
$m^6/(\text{kmole})^2$	c_1	+0.86687
	c_2	-0.42191
	c_3	+0.10118
	c_4	-0.00958
$d_i \cdot 10^2$	d_0	+0.14604
$m^9/(\text{kmole})^3$	d_1	-0.11465
	d_2	+0.028960
	d_3	-0.002310
	d_4	0.00000
$e_i \cdot 10^3$	e_0	-0.14232
$m^{12}/(\text{kmole})^4$	e_1	+0.12606
	e_2	-0.035740
	e_3	+0.00325
	e_4	+0.00002

The dependence of the second virial coefficient B_{12} on temperature for a mixture is approximated by the polynomial

$$B_{12}(T) = \sum_{i=0}^3 b_i T^i, \text{ cm}^3/\text{mole}. \quad (10)$$

The coefficients of b_i were determined on a digital computer by the method of least squares for the whole aggregate of available experimental data [15-18] with respect to $B_{12}(T)$, which are given in Table 4. For approximating $C_{112}(T)$, $C_{122}(T)$, $\gamma_1(T)$, $\gamma_2(T)$, $\alpha_1(T)$ a polynomial for degrees of $1/T$ was selected

$$\begin{aligned} C_{112}(T) &= \sum_{i=0}^3 (c_i/T^i), \text{ m}^6/(\text{kmole})^2, \\ C_{122}(T) &= \sum_{i=0}^3 (c'_i/T^i), \text{ m}^6/(\text{kmole})^2, \\ \gamma_1(T) &= \sum_{i=0}^3 (d_i/T^i), \text{ m}^9/(\text{kmole})^3, \\ \gamma_2(T) &= \sum_{i=0}^3 (d'_i/T^i), \text{ m}^9/(\text{kmole})^3, \\ \alpha_1(T) &= \sum_{i=0}^3 (e_i/T^i), \text{ m}^{12}/(\text{kmole})^4. \end{aligned} \quad (11)$$

The coefficients of these dependences were determined on a digital computer by the method of least squares in three variables for the whole aggregate of available experimental data with respect to compressibility for a mixture of helium and nitrogen [2, 4, 15], also including our experimental data. The values of the coefficients are given in Table 5.

Table 4. Values of the second virial coefficient $B_{12}(T)$ for mixtures of helium and nitrogen according to the data of various authors.

T, °K	273,15	298,15	303,15	373,15	448,15	523,15	598,15	673,15	748,15			
cm ³ /mole	21,63	22,08	21,83	22,49	22,82	22,92	21,41	21,72	22,20	21,20	20,33	21,27
Author	[17]	MGM	[16]	MGM	MGM	[18]	[18]	[18]	MGM	[18]	[18]	MGM

Table 5. Values of the coefficients of polynomials (11) in the equation of state for a helium-nitrogen mixture.

b_i	b_0	+6,87490
cm^3/mole	b_1	+17,0850
	b_2	-5,87540
	b_3	+0,63099
$c_i \cdot 10^2$	c_0	+0,25786
$\text{m}^6/(\text{kmole})^2$	c_1	-0,85548
	c_2	+0,34218
	c_0'	-0,17026
	c_1'	+0,94896
	c_2'	-1,06670
$d_i \cdot 10^2$	d_0	+0,03981
$\text{m}^9/(\text{kmole})^3$	d_1	-0,42416
	d_2	+1,02320
	d_0'	+0,07720
	d_1'	-0,16915
	d_2'	-0,16507
$e_i \cdot 10^2$	e_0	-0,11255
$\text{m}^{12}/(\text{kmole})^4$	e_1	+0,72084
	e_2	-1,19520

The temperature functions for the pure components were calculated with equations (2), (3), (7). At a zero concentration of one of the components of the mixture the equation of state of the mixture (8) turns into the equation of state of the other pure component.

A total of 133 experimental points were taken for processing. The deviation of the calculated values of the coefficient of

compressibility from the experimental values for 76 points was less than $\pm 0.05\%$, for 37 points less than $\pm 0.10\%$, for 18 points less than $\pm 0.20\%$ and for only 2 points the deviation was more than 0.20% .

A comparison of the calculated values of the coefficients of compressibility and density of a mixture of helium and nitrogen, computed by equation of state (8), with the experimental values [2, 4, 15] shows a satisfactory correspondence in the entire temperature and pressure region, encompassed by the experiment. This makes it possible to expect, that in the whole temperature region $273.15-773.15^\circ\text{K}$ at pressures of up to 400 bar the proposed equation of state will make it possible to obtain reliable and accurate data for the caloric properties of a mixture of helium and nitrogen.

Conclusions. 1. The equations of state for helium, nitrogen and their mixtures describe the available experimental data for helium and nitrogen within limits of $\pm 0.01\%$, and for mixtures of helium and nitrogen $\pm 0.2\%$.

2. The values of the coefficients of compressibility and density in temperature interval $273.15-773.15^\circ\text{K}$ and pressure interval 50-400 bar, computed with the equation of state for a mixture of helium and nitrogen, can be used for calculating technological processes and solving planning and design problems.

BIBLIOGRAPHY

1. В. Н. Попов, В. И. Чернышов. Экспериментальное исследование сжимаемости смесей гелий — азот. I. Изв. АН СССР. Энергетика и транспорт, 1968, № 4.
2. В. Н. Попов, В. И. Чернышов. Экспериментальное исследование сжимаемости смесей гелий — азот. II. Изв. АН СССР. Энергетика и транспорт, 1968, № 5.
3. В. И. Чернышов, В. Н. Попов, Н. В. Цедерберг. Экспериментальное исследование сжимаемости гелия. Атомная энергия, 1969, т. 26, вып. 1.
4. В. Н. Попов, В. И. Чернышов. Экспериментальное исследование сжимаемости смесей азот — гелий. Теплоэнергетика, 1969, № 2.
5. A. Michels, U. Wouters. Isotherms of helium between 0 and 150°C up to 200 amagat. Physica, 1941, v. 8.

6. R. Wiebe, V. L. Gaddi, C. Heins. The compressibility isotherms of helium at temperatures from -70 to 200°C and at pressures to 1000 atmospheres. *J. Amer. Chem. Soc.*, 1931, v. 53, p. 1721.
7. L. Holborn, I. Otto. Ober die Isothermen einiger Gase zwischen $+400^{\circ}$ und -183° . *Z. Phys.*, 1925, v. 33, № 1.
8. C. W. Gabby, C. C. Tanner, J. Masson. The pressures of gaseous mixtures. II. Helium and hydrogen and their intermolecular forces. *Proc. Roy. Soc. A.*, 1929, v. 122.
9. W. H. Keesom, Y. Y. M. Van Santen. Isothermals of helium at temperatures of 0 , 20 and 100 degrees C, and pressures from 5.5 to 16.5 atmospheres. *Commun. Phys. lab. univers. leiden*, 1933, № 227 b.
10. W. G. Schneider. Compressibility of gases at high temperatures. *Canad. J. Res. B.* 1946, v. 27, p. 339.
11. W. G. Schneider, J. Duffie. Compressibility of gases at high temperatures. II. Second virial coefficients of helium in the temperature range 0 to 600°C . *J. Chem. Phys.*, 1949, v. 17, № 9.
12. Н. В. Цедерберг, В. Н. Попов, Н. А. Морозова. Теплофизические свойства гелия. Госэнергоиздат, 1961.
13. A. Michels, H. Wouters, J. de Boer. Isotherms of nitrogen between 0° and 150° at pressures from 20 to 80 atm. *Physica*, 1934, v. 1, p. 587.
14. J. Saurel. Appareillage pour la détermination des équations d'état des gaz comprimés aux températures élevées application à l'étude de l'azote jusqu'à 100 kg/cm^2 et 1000°C . *J. rech. Centra nat. rech. Sci.*, 1958, № 42.
15. R. J. Witonsky, J. G. Miller. Compressibility of gases. IV. The Burnett method applied to gas mixtures at higher temperatures. The second virial coefficients of the helium-nitrogen system from 175° to 475° . *J. Amer. Chem. Soc.*, 1963, v. 85, № 3.
16. D. A. Gyorog, E. F. Obert. Virial coefficients for argon, methane, nitrogen, xenon. *Amer. Inst. Chem. Engrs J.*, 1961, v. 10, № 5.
17. F. B. Canfield, T. W. Leland, R. Kobayashi. Compressibility factor for helium-nitrogen mixtures. *J. Chem. and Engng Data*, 1965, v. 10, № 2.
18. G. M. Kramer, J. G. Miller. Compressibility of gases. III. The second and third virial coefficients of mixtures of helium and nitrogen at 30°C . *J. Phys. Chem.*, 1957, v. 61, № 6.

Submitted 4 March 1970